

KEYTRUDA[®] (pembrolizumab)

May cause acute and serious side effects which can sometimes become life-threatening and lead to death. These may happen any time during treatment or even after your treatment has ended. You may experience more than one side effect at the same time.

Contact your doctor right away if you develop any side effects. This also applies to side effects that are not listed on this card.

Your doctor may give you other medicines in order to prevent more severe complications and reduce your symptoms.

Due to side effects, your doctor may consider it necessary to withhold the next dose of KEYTRUDA. In case of serious side effects, treatment with KEYTRUDA can be stopped.

For more information, consult the package leaflet for KEYTRUDA at www.felleskatalogen.no

IMPORTANT

- Do not be tempted to treat side effects yourself.
- Do not stop your treatment with KEYTRUDA unless you have discussed this with your doctor.
- **Take this card with you at all times**, especially when you travel, whenever you go to the emergency room, or when you must see another doctor.
- Be sure to notify any health care professional you see that you are being treated with KEYTRUDA and show them this card.

Patient Card

Lungs

- Shortness of breath
- Chest pain
- Coughing

Intestines

- Diarrhea or more bowel movements than usual
- Stools that are black, tarry, sticky, or contain blood or mucus
- Severe stomach pain or tenderness
- Nausea or vomiting

Liver

- Nausea or vomiting
- Feeling less hungry
- Pain on the right side of stomach
- Yellowing of skin or whites of eyes
- Dark urine
- Bleeding or bruising more easily than normal

Kidneys

- Changes in the amount or color of your urine

Hormone glands

- Rapid heartbeat
- Weight loss or weight gain
- Increased sweating
- Hair loss
- Feeling cold
- Constipation
- Deeper voice
- Muscle aches
- Dizziness or fainting
- Headaches that will not go away or unusual headache

Type 1 diabetes, including diabetic ketoacidosis

- Feeling more hungry or thirsty than usual
- Needing to urinate more often
- Weight loss
- Feeling tired or feeling sick
- Stomach pain
- Fast and deep breathing
- Confusion
- Unusual sleepiness
- A sweet smell to your breath
- A sweet or metallic taste in your mouth
- A different odour to your urine or sweat

• If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects electronically to the Norwegian Medical Products Agency: www.dmp.no/pasientmelding.
• By reporting side effects, you can help provide more information on the safety of this medicine.

Important Contact Information

Name of Doctor

Office Phone

After-hours Phone

My Name

My Phone

Emergency Contact (Name and Phone)

Important Information for Health Care Providers

This patient is being treated with KEYTRUDA (pembrolizumab), which can cause acute and serious immune-mediated adverse reactions that may appear any time during treatment or even after treatment. Assess the patient for signs and symptoms of immune-mediated adverse reactions. Early diagnosis and appropriate management are essential to minimise any consequences of immune-mediated adverse reactions.

For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm aetiology or exclude other causes. Specific guidelines for managing immune-mediated adverse reactions are available in the Summary of Product Characteristics (SPC) for KEYTRUDA.

Consult an oncologist or other medical specialist for management of organ-specific immune-mediated adverse reactions.

Please consult the Summary of Product Characteristics (SPC) for KEYTRUDA at www.felleskatalogen.no or contact MSD (Norge) AS, Medical Information, on +47 32 20 73 00 for more information.



MSD (Norge) AS, Pb. 1579 Vika, 0118 Oslo, tel. + 47 32 20 73 00.
Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates.
All rights reserved. NO-KEY-00516 10/24

Version 13, October 2024